

Topical review

Genital pain in women: Beyond interference with intercourse

Sophie Bergeron^{a,b,*}, Natalie O. Rosen^{a,c}, Mélanie Morin^d^a Department of Psychology, Université de Montréal, Montréal, QC, Canada^b Department of Psychology, McGill University Health Centre, Montréal, QC, Canada^c Department of Obstetrics and Gynecology, IWK Health Centre, Halifax, NS, Canada^d School of Rehabilitation, Université de Sherbrooke, Sherbrooke, QC, Canada

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

1. Introduction

Pelvic pain problems affecting women's sexual and reproductive health are poorly understood and often misdiagnosed or ignored [21]. In addition to their high prevalence rates, from 12% to 21% in community samples, these conditions can impinge on patients' sexual functioning, psychological well-being, and quality of life [1]. Recommendations for the classification of sexual pain in the upcoming *Diagnostic and Statistical Manual of Mental Disorders-V* suggest collapsing dyspareunia and vaginismus into a single diagnostic entity called *genito-pelvic pain/penetration disorder*, which comprises five elements: percentage success of vaginal penetration, pain with vaginal penetration, fear of vaginal penetration or of pain during vaginal penetration, pelvic floor muscle dysfunction, and medical comorbidity [6]. Noteworthy is the emphasis on the multidimensional aspects of sexual pain, which represents an improvement over previous classifications. Yet genital pain remains within the category of sexual dysfunction, despite accumulating evidence in favour of a pain conceptualization. Further, the diagnosis still focuses heavily on the interference of the pain with a single type of sexual activity, intercourse. These classification challenges are reflected in our search for etiological explanations, with proponents of biomedical causes on one side [8] and those adhering to psychosexual models on the other [17]. We propose an integrated model taking into account the interdependency of biomedical – including pelvic floor muscle dysfunctions – cognitive, affective, behavioural, and interpersonal factors in genital pain and associated sexual impairment. Specifically, this model acknowledges that the pain experienced by women goes beyond an interference with intercourse, affecting and being affected by other sexual and nonsexual aspects of their lives.

2. Biomedical factors

Up until recently, most biomedical hypotheses had been derived from poorly controlled studies. In the last 10 years however,

the quality of the research has greatly improved, although it has pertained almost exclusively to provoked vestibulodynia – a burning pain triggered by pressure applied to the vulvar vestibule. The following factors have been found to be more common in women with genital pain than in controls: early puberty and pain with first tampon use [23], a history of repeated yeast infections [3], early and prolonged use of oral contraceptives [11], polymorphisms in genes regulating inflammatory response [19], nociceptor proliferation and sensitization [9,10], and lower touch and pain thresholds in the vulvar area as well as other body locations [31]. Taken together, findings suggest that both peripheral and central mechanisms are at play in the onset and maintenance of genital pain. Pelvic floor muscle dysfunctions figure prominently among the mechanisms receiving increased empirical attention.

3. Pelvic floor muscle dysfunctions

Studies to date indicate that abnormality at rest is the dysfunction of the pelvic floor muscles (PFM) that seems of greatest importance in the pathophysiology of genital pain. Different terminologies have been used to describe this abnormality, namely overactivity [29], hypertonicity [34], spasm [33], and spasticity [25]. Such abnormality is hypothesized to close the vaginal hiatus and interfere with penetration [4]. PFM hypertonicity evaluated by digital palpation [33,34] as well as increases in resting electromyographic (EMG) activity [18] were found in women with vestibulodynia and vaginismus compared to controls. It has been suggested that women with dyspareunia may exhibit a protective-like defensive reaction of the PFM during attempted vaginal penetration [33,34]. Shafik and El-Sibai [36] confirmed that reaction by showing greater resting EMG activity in women with vaginismus during dilator insertion compared to asymptomatic women.

Some studies have also shown that women with dyspareunia present deficits in the PFM contractile capacities. Reductions in strength and endurance were observed in women with dyspareunia compared to controls using palpation [33,34] and EMG [20,33]. However, measurement issues limit the validity of these findings, as EMG is not suitable for inter-subject comparison [2] and digital palpation is criticized for being a subjective measure [7]. New methodologies such as dynamometry and ultrasound appear more promising [27,30].

* Corresponding author. Address: Département de psychologie, Université de Montréal, C.P. 6128, Succursale Centre-Ville, Montréal, QC, Canada H3C 3J7. Tel.: +1 514 343 6111x5353; fax: +1 514 343 2285.

E-mail address: sophie.bergeron.2@umontreal.ca (S. Bergeron).

Whether PFM dysfunctions are a consequence or a cause of genital pain remains unknown. It has been hypothesized that genital pain and inflammation may provoke PFM dysfunctions by (1) destabilizing the PFM, inducing tensions and poor muscle control [20]; (2) triggering a defensive-like reaction toward pain during penetration attempts [33,34]; (3) intruding the underlying muscle tissues and causing hypersensitivity in the PFM [39]. Others have suggested that PFM dysfunctions may initiate an increase in mucosal sensitivity [40]. It is likely that a vicious cycle involving pain and further muscle dysfunctions is implicated, potentially compounded by the involvement of cognitive, affective, and behavioural factors.

4. Cognitive, affective, and behavioural factors

Research published in the last decade indicates that psychosocial factors may contribute to the onset of genital pain problems. In a study involving over 1400 adolescent girls, those complaining of dyspareunia of at least 6 months' duration were more likely to report past sexual abuse, fear of physical abuse, and trait anxiety in comparison to no-pain controls [26]. Additionally, the first study focusing on victimization in a general population adult sample showed that severe physical or sexual childhood abuse was linked to a 4- to 6-fold risk of reporting genital pain in adulthood [22]. These findings need not be interpreted solely in light of a sexual development perspective. Longitudinal work following individuals from birth to 32 years has shown that exposure to childhood maltreatment is linked to clinically significant biomarkers of inflammation in adulthood such as C-reactive protein levels, suggesting that inflammation may mediate the effect of adverse early life experiences on adult health [14]. Further, in a randomized treatment outcome study, pretreatment levels of erotophobia – the tendency to respond with negative affect to sexual cues – were found to account for 27% of the variance in a surgical outcome for vestibulodynia, independent of pretreatment pain [5]. Taken together, these results indicate that psychosocial factors may contribute to the development and persistence of genital pain.

Contrary to clinical lore, pain and sexual impairment may be distinct and partially independent phenomena, as shown by the lack of significant correlation between pain and sexual function, as well as pain and sexual satisfaction, in a recent study focusing on women with vestibulodynia [35]. It may be more fruitful to search for different predictors of these two critical outcomes. One such study indicated that higher levels of hypervigilance, fear of pain, and catastrophizing were significant predictors of increased pain in this population, whereas greater anxiety and avoidance were associated with poorer sexual function. However, results also showed that lower levels of self-efficacy were related to worse pain and sexual function, suggesting that some variables may affect both outcomes indiscriminately [15].

Because research examining the relation between psychosocial factors and genital pain is still in its infancy, the majority of studies have been descriptive and cross-sectional. Such designs do not elucidate the extent to which pain intensity may drive patients to fear and avoid threat or whether fear and anxiety precede the experience of pain. The fear-avoidance model – whereby an initial pain experience possibly caused by an injury may be interpreted as threatening (catastrophizing), leading to fear of pain and to avoidant behaviours, which in turn lead to hypervigilance followed by disability (in this case, sexual dysfunction), disuse (potential reduction of the sexual repertoire), and depression – may provide a useful framework for integrating findings concerning muscular reactivity and psychological factors [38], especially when expanding the focus from intraindividual to interpersonal variables.

5. Interpersonal factors

Surprisingly, relationship variables have been widely neglected in dyspareunia and vaginismus. Although there appear to be no differences in self-reported dyadic adjustment [28] and no association between dyadic adjustment and pain in women with dyspareunia [16], interpersonal factors remain relevant given that partners both elicit and witness women's pain experience. Recent studies have identified key relational variables, including attachment orientation, attributions, and partner responses that underscore the need for further research in this area [16,24].

Communicating pain to a significant other may serve to evoke empathic responses, assistance, or to maximize proximity to one's partner [37]. In turn, partner responses to expressions of pain may reinforce and perpetuate patients' pain experience [13,35]. In women with vestibulodynia, higher solicitous partner responses were associated with greater pain during intercourse [16,35]. Partner solicitousness may encourage avoidance of sexual activity and enhance negative cognitive-affective factors, which in turn heighten pain. Greater solicitous and lower negative partner responses were also associated with more sexual satisfaction in women [35]. Partner sensitivity to women's pain may impact sexual satisfaction because couples may be more or less likely to focus on alternative sexual activities to penetration or on the emotional benefits of sexual activity [35].

What type of partner response will minimize women's pain intensity, but maximize her sexual satisfaction? Facilitative responses, in which the partner encourages the patient's efforts at coping with the pain, warrant further investigation as they are associated with less patient disability in chronic pain populations [32]. In women with genital pain, facilitative responses may promote engaging in sexual activity by focusing on less painful activities or by expressing affection and pleasure during or after sexual activity.

Recent calls in the literature have lobbied for moving beyond cognitive-behavioural models of pain interactions in couples to include intimacy models [12]. Cognitive-behavioural models are limited in their focus on pain ratings and descriptions of the pain, often failing to capture pain-related distress, which can be another facet of verbal pain behaviours [12]. In contrast, intimacy models emphasize emotional disclosure and subsequent validating or invalidating partner responses. While cognitive-behavioural models view partner responses as reinforcing pain behaviours, intimacy models assert that these may enhance or disrupt intimacy and emotion regulation in couples [12]. It may be that the association between women's genital pain and disclosure about the pain will vary according to the partner's level of emotional validation, which in turn may contribute to pain persistence. Researchers will benefit from drawing on several models of interpersonal processes in order to better understand the role of dyadic interactions in couples with genital pain.

6. Conclusion

Fragmentation of scientists into specific disciplines, ongoing classification issues, and lack of sound research designs all impede progress concerning the identification of etiologic factors and the development of tailored interventions for women with genital pain. Empirical evidence to date suggests the existence of multiple etiologic pathways leading to the development and persistence of genital pain. We suggest that initial biomedical/mechanical trauma to the genitalia may trigger inflammatory processes, pelvic floor muscle dysfunctions, and other local changes leading to nociceptor sensitization and further peripheral and central alterations in pain processing. The experience of pain combined with a lack of proper diagnosis and treatment by health professionals [21] may interact

with individual predispositions and generate varying degrees of psychological distress in patients and their partners. In turn, cognitive, behavioural, affective, and interpersonal factors may modulate pain intensity as well as associated sexual impairment, as per the fear-avoidance model [38], such that not all women who have an initial experience of pain are at risk of suffering from a persistent condition, or of developing sexual dysfunction. Further, predictors of the pain experience can be different from those of the negative sexual and psychological symptomatology. Genital pain may be best conceptualized as a multidimensional persistent pain condition that constitutes the endpoint of multiple etiologic trajectories and whose understanding goes beyond its interference with intercourse.

Further progress in the genital pain field demands an expansion of our research methodologies. First, future research should include both members of the couples in order to examine the dynamic interactions of interpersonal processes on pain and associated psychological distress and sexual impairment. Second, it is important to venture beyond the study of clinical samples, which are probably biased in the direction of increased symptomatology. Third, researchers should use prospective and experimental designs such as observational and experience processing methods to tease apart the temporal order of the associations among women's behaviours, partner responses, pain, and psychosexual variables. Fourth, biomedical factors such as pelvic floor muscle dysfunctions should be investigated using better validated measurements to understand further their implication in the pathophysiology of genital pain. Such studies will contribute to the development of more refined, biopsychosocial etiological models of genital pain and will inform the development of targeted interventions.

Conflict of interest statement

None of the authors has a conflict of interest.

Acknowledgments

This work was supported by Canadian Institutes of Health Research (CIHR) and Social Sciences and Humanities Research Council grants awarded to Sophie Bergeron and by CIHR postdoctoral fellowships awarded to Natalie O. Rosen and Mélanie Morin.

References

- Arnold LD, Bachmann GA, Rosen R, Kelly S, Rhoads GG. Vulvodynia: characteristics and associations with comorbidities and quality of life. *Obstet Gynecol* 2006;107:617–24.
- Auchincloss CC, McLean L. The reliability of surface EMG recorded from the pelvic floor muscles. *J Neurosci Methods* 2009;182:85–96.
- Bachmann GA, Rosen R, Arnold LD, Burd I, Rhoads GG, Leiblum SR, Avis N. Chronic vulvar and other gynecologic pain: prevalence and characteristics in a self-reported survey. *J Reprod Med* 2006;51:3–9.
- Basson R, Berman J, Burnett A, Derogatis L, Ferguson D, Fourcroy J, Goldstein I, Graziottin A, Heiman J, Laan E, Leiblum S, Padma-Nathan H, Rosen R, Seagraves K, Seagraves RT, Shabsigh R, Sipski M, Wagner G, Whipple B. Report of the international consensus development conference on female sexual dysfunction: definitions and classifications. *J Urol* 2000;163:888–93.
- Bergeron S, Khalifé S, Glazer HI, Binik YM. Surgical and behavioral treatments for vestibulodynia: two-and-one-half year follow-up and predictors of outcome. *Obstet Gynecol* 2008;111:159–66.
- Binik Y. The DSM diagnostic criteria for dyspareunia. *Arch Sex Behav* 2010;39:292–303.
- Bo K, Finckenhagen HB. Vaginal palpation of pelvic floor muscle strength: inter-test reproducibility and comparison between palpation and vaginal squeeze pressure. *Acta Obstet Gynecol Scand* 2001;80:883–7.
- Bohm-Starke N. Medical and physical predictors of localized provoked vulvodynia. *Acta Obstet Gynecol Scand* 2010;89:1504–10.
- Bohm-Starke N, Hilliges M, Brodda-Jansen G, Rylander E, Torebjork E. Psychophysical evidence of nociceptor sensitization in vulvar vestibulitis syndrome. *Pain* 2001;94:177–83.
- Bohm-Starke N, Hilliges M, Falconer C, Rylander E. Neurochemical characterization of the vestibular nerves in women with vulvar vestibulitis syndrome. *Gynecol Obstet Invest* 1999;48:270–5.
- Bouchard C, Brisson J, Fortier M, Morin C, Blanchette C. Use of oral contraceptive pills and vulvar vestibulitis: a case-control study. *Am J Epidemiol* 2002;156:254–61.
- Cano A, de C Williams AC. Social interaction in pain: reinforcing pain behaviors or building intimacy? *Pain* 2010;149:9–11.
- Cano A, Johansen AB, Leonard MT, Hanawalt JD. What are the marital problems of patients with chronic pain? *Curr Pain Headache Rep* 2005;9:96–100.
- Danese A, Pariante CM, Caspi A, Taylor A, Poulton R. Childhood maltreatment predicts adult inflammation in a life-course study. *Proc Natl Acad Sci USA* 2007;104:1319–24.
- Desrochers G, Bergeron S, Khalifé S, Dupuis MJ, Jodoin M. Fear avoidance and self-efficacy in relation to pain and sexual impairment in women with provoked vestibulodynia. *Clin J Pain* 2009;25:520–7.
- Desrosiers M, Bergeron S, Meana M, Leclerc B, Binik YM, Khalife S. Psychosexual characteristics of vestibulodynia couples: partner solicitousness and hostility are associated with pain. *J Sex Med* 2008;5:418–27.
- Dewitte M, Van Lankveld J, Crombez G. Understanding sexual pain: a cognitive-motivational account. *Pain* 2011;152:251–3.
- Frasson E, Graziottin A, Priori A, Dall'ora E, Didonè G, Garbin EL, Vicentini S, Bertolasi L. Central nervous system abnormalities in vaginismus. *Clin Neurophysiol* 2009;120:117–22.
- Gerber S, Bongiovanni AM, Ledger WJ, Witkin SS. Defective regulation of the proinflammatory immune response in women with vulvar vestibulitis syndrome. *Am J Obstet Gynecol* 2002;186:696–700.
- Glazer HI, Jantos M, Hartmann EH, Swencionis C. Electromyographic comparisons of the pelvic floor in women with dysesthetic vulvodynia and asymptomatic women. *J Reprod Med* 1998;43:959–62.
- Harlow BL, Stewart EG. A population-based assessment of chronic unexplained vulvar pain: have we underestimated the prevalence of vulvodynia? *J Am Med Womens Assoc* 2003;58:82–8.
- Harlow BL, Stewart EG. Adult-onset vulvodynia in relation to childhood violence victimization. *Am J Epidemiol* 2005;161:871–80.
- Harlow BL, Wise LA, Stewart EG. Prevalence and predictors of chronic lower genital tract discomfort. *Am J Obstet Gynecol* 2001;185:545–50.
- Jodoin M, Bergeron S, Khalifé S, Dupuis MJ, Desrochers G, Leclerc B. Male partners of women with provoked vestibulodynia: attributions for pain and their implications for dyadic adjustment, sexual satisfaction, and psychological distress. *J Sex Med* 2008;5:2862–70.
- Kuijpers HC, Bleijenberg G. The spastic pelvic floor syndrome. A cause of constipation. *Dis Colon Rectum* 1985;28:669–72.
- Landry T, Bergeron S. Biopsychosocial factors associated with dyspareunia in a community sample of adolescent girls. *Arch Sex Behav* 2010 [Epub ahead of print].
- Majida M, Braekken IH, Umek W, Bo K, Saltyte Benth J, Ellstrom Engh M. Interobserver repeatability of three- and four-dimensional transperineal ultrasound assessment of pelvic floor muscle anatomy and function. *Ultrasound Obstet Gynecol* 2009;33:567–73.
- Meana M, Binik YM, Khalife S, Cohen DR. Biopsychosocial profile of women with dyspareunia. *Obstet Gynecol* 1997;90:583–9.
- Messelink B, Benson T, Berghmans B, Bo K, Corcos J, Fowler C, Laycock J, Lim PH, van Lunsen R, á Nijeholt GL, Pemberton J, Wang A, Watier A, Van Kerrebroeck P. Standardization of terminology of pelvic floor muscle function and dysfunction: report from the pelvic floor clinical assessment group of the International Continence Society. *Neurourol Urodyn* 2005;24:374–80.
- Morin M, Gravel D, Bourbonnais D, Dumoulin C, Ouellet S. Reliability of dynamometric passive properties of the pelvic floor muscles in postmenopausal women with stress urinary incontinence. *Neurourol Urodyn* 2008;27:819–25.
- Pukall CF, Binik YM, Khalifé S, Amsel R, Abbott FV. Vestibular tactile and pain thresholds in women with vulvar vestibulitis syndrome. *Pain* 2002;96:163–75.
- Raichle KA, Romano JM, Jensen MP. Partner responses to patient pain and well behaviors and their relationship to patient pain behavior, functioning, and depression. *Pain* 2011;152:82–8.
- Reissing ED, Binik YM, Khalifé S, Cohen D, Amsel R. Vaginal spasm, pain, and behavior: an empirical investigation of the diagnosis of vaginismus. *Arch Sex Behav* 2004;33:5–17.
- Reissing ED, Brown C, Lord MJ, Binik YM, Khalifé S. Pelvic floor muscle functioning in women with vulvar vestibulitis syndrome. *J Psychosom Obstet Gynaecol* 2005;26:107–13.
- Rosen NO, Bergeron S, Leclerc B, Lambert B, Steben M. Woman and partner-perceived partner responses predict pain and sexual satisfaction in provoked vestibulodynia (PVD) couples. *J Sex Med* 2010;7:3715–24.
- Shafik A, El-Sibai O. Study of the pelvic floor muscles in vaginismus: a concept of pathogenesis. *Eur J Obstet Gynecol Reprod Biol* 2002;105:67–70.
- Sullivan MJ, Thorn B, Haythornthwaite JA, Keefe F, Martin M, Bradley LA, Lefebvre JC. Theoretical perspectives on the relation between catastrophizing and pain. *Clin J Pain* 2001;17:52–64.
- Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000;85:317–32.
- Wesselmann U. Neurogenic inflammation and chronic pelvic pain. *World J Urol* 2001;19:180–5.
- Zolnoun D, Hartmann K, Lamvu G, As-Sanie S, Maixner W, Steege J. A conceptual model for the pathophysiology of vulvar vestibulitis syndrome. *Obstet Gynecol Surv* 2006;61:395–401.